

Contents

Abbreviations	XI
Introduction	1
1. Historical overview	1
2. Synthesis, release and catabolism of ATP	2
2.1 Vesicular release of ATP	3
2.2 Anion channel-mediated release of ATP	4
2.3 Connexin hemichannel-mediated ATP release	4
3. Synthesis, release and catabolism of adenosine.	6
3.1 Adenosine release	7
3.2 Synthesis of adenosine at extracellular level	8
Purinergic receptors	9
1. P1 receptors	9
2. Pharmacology of P1 receptors	10
3. Distribution of P1 receptors.	11
3.1 Central Nervous System (CNS)	12
4. P2 receptors	13
5. P2X receptors	14
6. Pharmacology of P2X receptors	18
6.1 Agonists	18
6.2 Antagonists	19
7. P2Y receptors	20
8. Pharmacology of P2Y receptors.	22
8.1 Agonists	22
8.2 Antagonists	23
9. Distribution of P2 receptors	24
9.1 Neuronal localization of P2X receptors	26
9.2 Neuronal localization of P2Y receptors	27
Effects of extracellular purines	29
1. Role of P1 receptors	29
1.1 P1 receptors and neurotransmission.	30
1.1 P1 receptors and cerebral ischemia	32
2. Anoxic depolarization	35

3.	Role of P2 receptors	38
3.1	P2 receptors and neurotransmission	39
3.2	P2 receptors and cerebral ischemia	43
3.3	P2 receptors and trophic effects	44

Aim of the study	46
-------------------------	----

Materials and methods	47
------------------------------	----

1.	Extracellular recordings	47
1.1	Acute rat hippocampal slices preparation.	47
1.2	Experimental procedure	47
1.3	Protocols used	48
1.4	OGD conditions	48
2.	Patch clamp recordings	49
3.	Cell cultures	50
4.	Solutions	51
5.	Patch pipettes fabrication	51
6.	Experimental procedures.	52
7.	Voltage clamp protocols	53
8.	Current clamp experiments.	54
9.	Measurement of ATP release from hMSCs	55
10.	Chemicals	55
11.	Statistical analysis	61

RESULTS. Section 1

ROLE OF PURINES IN CA1 HIPPOCAMPAL NEUROTRANSMISSION
UNDER NORMOXIC CONDITIONS AND DURING OGD. EXTRACELLULAR
RECORDINGS FROM RAT HIPPOCAMPAL SLICES

Chapter 1

Role of A₃ adenosine receptors during OGD	65
---	----

1.	Historical background	65
2.	Results	66
2.1	Experiments were performed on a total of 219 slices taken from 133 rats	66
3.	Discussion	78
3.1	Role of adenosine receptor antagonists in depression and disruption of CA1 hippocampal excitatory synaptic transmission during <i>in vitro</i> ischemia.	78
3.2	Therapeutic implications for the protective role of adenosine A ₃ receptor antagonists during prolonged ischemic conditions.	81
3.3	Role of adenosine receptor agonists in depression and disruption of CA1 hippocampal excitatory synaptic transmission during <i>in vitro</i> ischemia.	82
3.5	Conclusions	83

Chapter 2

Role of P2 purinergic receptors on hippocampal CA1 neurotransmission under normoxic conditions and during OGD

	85
1. Historical background	85
2. Results	86
2.1 Effects of ATP on CA1 hippocampal neurotransmission	86
2.2 Effects of ATP γ S on CA1 hippocampal neurotransmission	90
2.3 Effects of ATP and ATP γ S on PPF in the CA1 region of rat hippocampus	93
2.4 Effects of different purinergic antagonists on OGD-evoked disruption of neurotransmission in the CA1 region of rat hippocampal slices	94
3. Discussion	97
3.1 Role of P2 receptors in hippocampal neurotransmission under normoxic conditions	97
3.2 Role of P2 receptors in hippocampal neurotransmission during OGD	98
3.3 Conclusions	99

RESULTS. Section 2

STUDIES ON P2 RECEPTOR MODULATION OF MEMBRANE IONIC CURRENTS BY PATCH CLAMP RECORDINGS

Chapter 3

Role of P2Y₁ receptor stimulation in medium spiny neurons of rat striatal slices

	103
1. Historical background	103
2. Results	104
2.1 P2Y ₁ receptor activation increases the outward currents elicited by a voltage ramp depolarization in striatal medium spiny neurons	104
2.2 Characterization of the P2Y ₁ -activated conductance in striatal medium spiny neurones.	106
2.3 Single channel recordings show SK potassium channel opening following P2Y ₁ receptor activation	109
2.4 2MeS-ADP decreases the firing rate of medium spiny neurons recorded under the current-clamp mode in striatal slices	110
2.5 Spontaneous miniature synaptic currents are increased in amplitude during the application of 2meS-ADP.	112
3. Discussion	113

Chapter 4

Role of purine nucleotides in cultured hMSCs: release of ATP and activation of P2 receptors

	119
1. Historical background	119
2. Results	120
2.1 Electrophysiological experiments	121
2.2 Effects of exogenous ATP in potassium-free conditions	124

2.3	Effects of ATP in hMSCs in the perforated patch-clamp configuration	126
3.	Discussion	128
3.1	Conclusion	130
Chapter 5		
Pharmacological characterization of the recently deorphanized GPR17 receptor transfected in 1321N1 astrocytoma cells.		
1.	Historical background	131
2.	Results	132
2.1	Exogenously applied UDP-glucose and UDP-galactose activate outward potassium currents in 1321N1 astrocytoma cell stably transfected with GPR17.	132
2.2	GPR17 receptor, stably transfected 1321N1 astrocytoma cells, is also sensitive to leukotriene-related compounds	134
3.	Discussion	137
Final remarks		141
References		145